



ONDERSTPOORT
BIOLOGICAL PRODUCTS

For animal use only

ANTHRAX SPORE VACCINE

Reg. No. G 0112 (Act 36/1947) Namibia: V01 /24.4/138

Glycerine suspension of avirulent *Bacillus anthracis* 34F2 spores for the immunisation of cattle, horses, mules, donkeys, sheep, goats, pigs and camels against anthrax.

Store the vaccine at a temperature of 4 °C to 8 °C, in a refrigerator. Do not use after the expiry date printed on the bottle.

RECOMMENDATIONS FOR USE

All animals should be inoculated annually. Working animals should be rested for a week to 10 days after inoculation. Until the swelling has subsided, these animals should not be driven over long distances.

WARNINGS

Do not slaughter animals for human consumption within 21 days of vaccination. Vaccinate healthy animals only. Keep out of reach of children, uninformed persons and animals. Goats must be injected in the inner thigh or tailfold as these animals are inclined to show allergic reactions with swellings at the site of injection, which may cause suffocation if the inoculation is given in the neck. Despite all possible precautionary measures, the absolute safety of this vaccine can unfortunately not be guaranteed in all animals. Occasionally animals, on certain farms, may show shock reactions shortly after injection with the vaccine. Users of the vaccine are, therefore, advised to first inject a number of young animals and observe them for 30-60 minutes. Should they show no adverse reactions, the rest of the herd may be immunised. Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE

Use only as directed.

Sterilise syringes and needles by boiling in water for at least 15 minutes. Do not use disinfectants or methylated spirits for sterilising either syringes or needles. Use a separate needle for each animal. Shake the bottle well before use. Avoid exposure to high temperatures and direct sunlight during inoculation.

DOSAGE: 1 ml subcutaneously for all animals irrespective of species, breed or age.

EFFECTS OF THE VACCINE

The vaccine causes a swelling at the site of injection. The swelling is temporary and subsides after a few days. Immunity develops within 2-4 weeks after vaccination and persists for approximately 9-12 months but cannot be guaranteed in all animals.

PACKING

Available in bottles of 50 and 100 doses.

Registration holder:

Onderstepoort Biological Products SOC Ltd, Co. Reg. No. 2000/022686/06
Private Bag X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Fax: +27 (0) 12 522 1591

Made in South Africa

Edition 6

Lomax Printers
WT 17220

P1015



ONDERSTAPOORT
BIOLOGIESE PRODUKTE

Slegs vir dieregebruik

MILTSIEKTE SPOOR ENTSTOF

Reg. Nr. G 0112 (Wet 36/1947) Namibië: V01 /24.4/138

Gliserien suspensie van avirulente *Bacillus anthracis* 34F2 spore vir die immunisering van beeste, perde, muile, donkies, skape, bokke, varke en kamele teen miltsiekte.

Bewaar die entstof by 'n temperatuur van 4 °C tot 8 °C, in 'n yskas. Moet nie die entstof na die vervaldatum wat op die bottel gedruk is gebruik nie.

AANBEVELINGS VIR GEBRUIK

Ent alle diere jaarliks. Diere wat werk moet 'n week tot 10 dae rus nadat hulle ingeënt is. Moet nie diere oor lang afstande aanjaag voordat die swelsel verdwyn het nie.

WAARSKUWINGS

Moet nie diere binne 21 dae na inenting vir menslike verbruik slag nie. Slegs gesonde diere moet ingeënt word. Hou buite bereik van kinders, ongeligte persone en diere. Bokke moet in die binnebond of stertvou ingeënt word, aangesien die diere geneig is om soms swaar allergiese reaksies met swelsels te ontwikkel op die plek van inspuiting en wat versmoring kan veroorsaak indien hulle in die nek ingeënt word. Ten spyte van alle moontlike voorsorgmaatreëls kan die absolute veiligheid van hierdie entstof ongelukkig nie in alle diere gewaarborg word nie. Enkele diere, op bepaalde plase, mag na toediening van die entstof skokreaksies toon. U word dus versoek om, wanneer entstof gebruik word, eers 'n tiental jong diere in te ent en vir 30-60 minute dop te hou. Indien hulle geen reaksie toon nie kan daar met die inenting van die res van die kudde voortgegaan word. Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS

Gebruik slegs soos voorgeskryf.

Steriliseer spuite en naalde deur dit vir ten minste 15 minute in water te kook. Moet nie ontsmettingsmiddels of brandspiritus gebruik om spuite en naalde te steriliseer nie. Gebruik 'n afsonderlike steriele naald vir elke dier. Skud die bottel goed voor gebruik. Vermy blootstelling aan hoë temperature en direkte sonlig gedurende inenting.

DOSIS: 1 ml onderhuids ongeag van soort, ras of ouderdom.

UITWERKING VAN DIE ENTSTOF

'n Plaaslike swelsel ontwikkel op die plek van inspuiting. Die swelsel is egter tydelik van aard en verdwyn na 'n paar dae. Immuniteit ontwikkel binne 2-4 weke na inenting en duur vir 9-12 maande maar kan egter nie in alle diere gewaarborg word nie.

VERPAKKING

Beskikbaar in bottels van 50 en 100 dosisse.

Registrasiehouer:

Onderstepoort Biologiese Produkte MSB Bpk, Mpy. Reg. Nr. 2000/022686/06
Privaatsak X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Faks: +27 (0) 12 522 1591

Vervaardig in Suid-Afrika

Weergawe 6

P1015