



ONDERSTPOORT
BIOLOGICAL PRODUCTS

For animal use only

LAMB DYSENTERY VACCINE

Reg. No. G 0102 (Act 36/1947) Namibia: NSR 0585

Formalinised alum-precipitated toxoid of *Clostridium perfringens (welchii)* type B for the immunisation of ewes and lambs against lamb dysentery.

Store the vaccine at a temperature of 4 °C to 8 °C, in a refrigerator. Do not use the vaccine after the expiry date printed on the bottle.

RECOMMENDATIONS FOR USE

Pregnant ewes are immunised because lambs contract the disease soon after birth. The first injection is given 4 weeks before the ewes are due to lamb and the second injection 3 weeks later. Ewes which have previously been injected need only receive 1 booster injection 10–14 days before lambing.

WARNINGS

Do not slaughter animals for human consumption within 21 days of vaccination. Vaccinate healthy animals only. Keep out of reach of children, uninformed persons and animals. Handle pregnant ewes with care. Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE

Use only as directed.

Sterilise syringes and needles by boiling in water for at least 15 minutes. Do not use disinfectants or methylated spirits for sterilising either syringes or needles. If possible use a sterile needle for each animal. Shake the bottle well before use. Avoid exposure to high temperature and direct sunlight during inoculation. Administer the vaccine subcutaneously in the groin or behind the shoulder.

DOSAGE: 2 ml subcutaneously

EFFECTS OF THE VACCINE

Other than a slight transient lameness in the leg nearest to the site of injection and a small hard nodule under the skin, no reaction is anticipated. Immunity is passed from the ewe to the lamb by means of the colostrum but cannot be guaranteed in all animals.

PACKING

Available in bottles of 50 doses.

Registration holder:

Onderstepoort Biological Products SOC Ltd, Co. Reg. No. 2000/022686/06
Private Bag X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Fax: +27 (0) 12 522 1591

Made in South Africa

Edition 4

Lomar Printers

P1161



ONDERSTEOORT
BIOLOGIESE PRODUKTE

Slegs vir dieregebruik

BLOEDPENS **ENTSTOF**

Reg. Nr. G 0102 (Wet 36/1947) Namibië: NSR 0585

Geformaliseerde aluin-gepresipiteerde toksoïed van *Clostridium perfringens (welchii)* tipe B vir die immunisering van ooie en lammers teen bloedpens.

Bewaar die entstof by 'n temperatuur van 4 °C tot 8 °C, in 'n yskas. Moet nie die entstof na die vervaldatum wat op die bottel gedruk is gebruik nie.

AANBEVELINGS VIR GEBRUIK

Omdat lammers die siekte kort na geboorte opdoen word die ooie ingeënt. Die eerste inenting vind plaas 4 weke voordat die ooie moet lam en die tweede inenting 3 weke later. Ooie wat voorheen ingeënt was ontvang slegs een skragingsdosis 10–14 dae voordat hulle moet lam.

WAARSKUWINGS

Moet nie diere binne 21 dae na inenting vir menslike verbruik slag nie. Slegs gesonde diere moet ingeënt word. Hou buite bereik van kinders, oningeligte persone en diere. Hanteer dragtige ooie met sorg. Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GEBRUIKSAANWYSINGS

Gebruik slegs soos voorgeskryf.

Steriliseer spuite en naalde deur dit vir ten minste 15 minute in water te kook. Moet nie ontsmettingsmiddels of brandspiritus gebruik om spuite en naalde te steriliseer nie. Gebruik sover moontlik 'n steriele naald vir elke dier. Skud die bottel goed voor gebruik. Vermoed blootstelling aan hoë temperature en direkte sonlig gedurende inenting. Dien die entstof onderhuids toe in die lies of agter die blad.

DOSIS: 2 ml onderhuids

UITWERKING VAN DIE ENTSTOF

Behalwe vir 'n geringe en verbygaande mankheid in die been naaste aan die plek van inspuiting en 'n klein harde knoppie onder die vel, word geen reaksie verwag nie. Immuniteit word deur middel van die biesmelk van die ooi aan die lam oorgedra maar kan egter nie in alle diere gewaarborg word nie.

VERPAKKING

Beskikbaar in bottels van 50 dosisse.

Registrasiehouer:

Onderstepoort Biologiese Produkte MSB Bpk, Mpy. Reg. Nr. 2000/022686/06
Privaatsak X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Faks: +27 (0) 12 522 1591

Vervaardig in Suid-Afrika

Weergawe 4

P1161